



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

158. Proposed by G. H. HARVILL, A. M., Malakoff, Texas.

Show that a law of density for points in space may be assumed such that the joint mass of any two points which are electrical images of each other in respect to a given sphere may be constant, and that their centers of gravity shall lie on the surface of the sphere.

Remark by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

This problem is the same as problem 148, a solution of which is given in No. 2, Vol. X, pages 48 and 49.

DIOPHANTINE ANALYSIS.

105. Proposed by HARRY S. VANDIVER, Bala, Pa.

Prove that every factor of $a^{2m} + b^{2m}$ is of the form $1[\text{mod}.2^{m+1}]$ where a and b are prime to each other.

Note by the PROPOSER.

The enunciation of this proposition is corrected on page 207, Vol. IX, of the MONTHLY, and it may be stated in full as follows:

If a , b and m are real positive integers, a and b relatively prime, then every odd factor of $a^{2m} + b^{2m}$ has the form $1[\text{mod}.2^{m+1}]$.

This special arithmetical theorem was first given by Euler, and may be directly proved by the principles in the theory of power residues. It is interesting, however, to examine its connection with a more general proposition. If $\omega_1, \omega_2, \dots, \omega_{\phi(n)}$ are all the distinct primitive roots of $x^n - 1 = 0$ and $\phi(n)$ the number of integers less than the positive integer n , and prime to it, then the rational function

$$F\{a, b, n\} = \{a - \omega_1 b\} \{a - \omega_2 b\} \dots \{a - \omega_{\phi(n)} b\}$$

possesses the following property:

Every divisor of $F\{a, b, n\}$ which is prime to n , has the form $1[\text{mod } n]$.

A proof of this for the case $b=1$, is to be found in Kronecker's *Vorlesungen über Zahlentheorie*, Vol. I, pages 440-441, and the analysis used may be readily extended to cover the case in which b is general. For $n=2^{m+1}$ (m , a positive integer) it will be noticed that

$$F\{a, b, 2^{m+1}\} = a^{2m} + b^{2m},$$

and by the general theorem above stated, it follows that every odd divisor of this function has the form $1[\text{mod } 2^{m+1}]$. When $a=2$, $b=1$, we may write

$$f\{m\} = 2^{2m} + 1,$$

and this form gives the remarkable numbers of Gauss which occur in the partition of the perigon.